

**Author Affiliations:** Institute for Translational Research in Children's Mental Health, University of Minnesota, Minneapolis (Patrick, Wagner); Institute of Child Development, University of Minnesota, Minneapolis (Patrick); Institute for Social Research, University of Michigan, Ann Arbor (Miech, Kloska, Johnston).

**Accepted for Publication:** October 2, 2019.

**Corresponding Author:** Megan E. Patrick, PhD, Institute for Translational Research in Children's Mental Health, University of Minnesota, 1100 Washington Ave S, Ste 101, Minneapolis, MN 55415 (mpatrick@umn.edu).

**Published Online:** April 6, 2020. doi:10.1001/jamapediatrics.2020.0175

**Author Contributions:** Dr Patrick and Ms Kloska had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

*Study concept and design:* Patrick.

*Acquisition, analysis, or interpretation of data:* All authors.

*Drafting of the manuscript:* Patrick, Kloska, Wagner.

*Critical revision of the manuscript for important intellectual content:* Patrick, Miech, Wagner, Johnston.

*Statistical analysis:* Patrick, Miech, Kloska.

*Obtained funding:* Patrick, Miech, Johnston.

*Administrative, technical, or material support:* Patrick, Miech, Johnston.

*Study supervision:* Patrick.

**Conflict of Interest Disclosures:** Dr Johnston has received grants from the US Department of Health and Human Services. No other disclosures were reported.

**Funding/Support:** Data collection and manuscript preparation were supported by research grant RO1DA001411 (Drs Miech and Johnston) and development of the manuscript was supported by research grant RO1DA037902 (Dr Patrick) from the National Institute on Drug Abuse.

**Role of the Funder/Sponsor:** The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Disclaimer:** The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute on Drug Abuse.

1. Johnston LD, Miech RA, O'Malley PM, Bachman JG, Schulenberg JE, Patrick ME. 2018 Overview: key findings on adolescent drug use. Accessed August 1, 2019. <http://www.monitoringthefuture.org/pubs/monographs/mtf-overview2018.pdf>
2. Johnson RM, Brooks-Russell A, Ma M, Fairman BJ, Tolliver RL Jr, Levinson AH. Usual modes of marijuana consumption among high school students in Colorado. *J Stud Alcohol Drugs*. 2016;77(4):580-588. doi:10.15288/jsad.2016.77.580
3. Mammen G, Rehm J, Rueda S. Vaporizing cannabis through e-cigarettes. *Can J Public Health*. 2016;107(3):e337-e338. doi:10.17269/CJPH.107.5747
4. Peters EN, Bae D, Barrington-Trimis JL, Jarvis BP, Leventhal AM. Prevalence and sociodemographic correlates of adolescent use and polyuse of combustible, vaporized, and edible cannabis products. *JAMA Netw Open*. 2018;1(5):e182765. doi:10.1001/jamanetworkopen.2018.2765
5. Friese B, Slater MD, Battle RS. Use of marijuana edibles by adolescents in California. *J Prim Prev*. 2017;38(3):279-294. doi:10.1007/s10935-017-0474-7
6. Miech RA, Johnston LD, O'Malley PM, Bachman JG, Schulenberg JE, Patrick ME. Volume I: secondary school students. Accessed August 1, 2019. [http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1\\_2018.pdf](http://www.monitoringthefuture.org/pubs/monographs/mtf-vol1_2018.pdf)

## Age-Related Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019 (COVID-19)

Children are susceptible to infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) but generally present with mild symptoms compared with adults.<sup>1</sup> Children drive spread of respiratory and gastrointestinal illnesses in the population,<sup>2</sup> but data on children as sources of SARS-CoV-2 spread are sparse.

Early reports did not find strong evidence of children as major contributors to SARS-CoV-2 spread,<sup>3</sup> but school clo-


sure early in pandemic responses thwarted larger-scale investigations of schools as a source of community transmission. As public health systems look to reopen schools and day cares, understanding transmission potential in children will be important to guide public health measures. Here, we report that replication of SARS-CoV-2 in older children leads to similar levels of viral nucleic acid as adults, but significantly greater amounts of viral nucleic acid are detected in children younger than 5 years.

**Methods** | Between March 23 and April 27, 2020, we performed SARS-CoV-2 reverse transcriptase-polymerase chain reaction (PCR) on nasopharyngeal swabs collected at various inpatient, outpatient, emergency department, and drive-through testing sites at a pediatric tertiary medical center in Chicago, Illinois. The Ann & Robert H. Lurie Children's Hospital of Chicago Institutional Review Board provided an exemption and full waiver of HIPAA authorization and informed consent. A Clinical Laboratory Improvement Amendments-certified laboratory analyzed samples using a US Food and Drug Administration Emergency Use Authorization PCR assay (Abbot RealTime SARS-CoV-2 Assay performed on the m2000 RealTime System [Abbott Laboratories]). PCR amplification cycle threshold (CT) values were recorded, with lower values indicating higher amounts of viral nucleic acid.

This cohort included all individuals aged younger than 1 month to 65 years who tested positive for SARS-CoV-2. Patients with symptoms suggestive of a COVID-19-compatible illness and/or high-risk exposures were tested. We included the first sample tested for patients with multiple samples. Because patients with severe infection have lower CT values,<sup>4</sup> we excluded 7 children who required supplemental oxygen support. We also excluded 7 asymptomatic patients, 29 patients with unknown duration of symptoms, and 19 patients whose symptoms started more than 1 week prior to testing. Swabs were collected using a standard bilateral nasopharyngeal sampling procedure. Several controls, including samples with known copy numbers, were included in each PCR run. Median and interquartile ranges for each group were measured and compared using the nonparametric Wilcoxon rank sum test. Two-sided *P* values less than .05 were considered statistically significant. Analyses were performed using Stata/IC statistical software version 16.0 (StataCorp).

**Results** | Our final cohort included 145 patients with mild to moderate illness within 1 week of symptom onset. We compared 3 groups: young children younger than 5 years (*n* = 46), older children aged 5 to 17 years (*n* = 51), and adults aged 18 to 65 years (*n* = 48). We found similar median (interquartile range) CT values for older children (11.1 [6.3-15.7]) and adults (11.0 [6.9-17.5]). However, young children had significantly lower median (interquartile range) CT values (6.5 [4.8-12.0]), indicating that young children have equivalent or more viral nucleic acid in their upper respiratory tract compared with older children and adults (Figure). The observed differences in median CT values between young children and adults approximate a 10-fold to 100-fold greater amount of SARS-CoV-2 in the upper respiratory tract of young children. We performed a sen-

**Figure. Distribution of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Reverse Transcriptase-Polymerase Chain Reaction (RT-PCR) Amplification Cycle Threshold (CT) Values From Nasopharyngeal Swabs Collected From Patients With Coronavirus Disease 2019**



Children younger than 5 years had significantly lower CT values compared with children aged 5 to 17 years ( $P = .02$ ) and adults 18 years and older ( $P = .001$ ). CT values were similar between children aged 5 to 17 years and adults 18 years and older ( $P = .34$ ). Midlines indicate the median, boxes indicate interquartile ranges, whiskers indicate the upper and lower adjacent values (within 1.5-fold the interquartile range), and isolated data points indicate outliers.

sitivity analysis and observed a similar statistical difference between groups when including those with unknown symptom duration. Additionally, we identified only a very weak correlation between symptom duration and CT in the overall cohort (Spearman  $\rho = 0.22$ ) and in each subgroup (young children, Spearman  $\rho = 0.20$ ; older children, Spearman  $\rho = 0.19$ ; and adults, Spearman  $\rho = 0.10$ ).

**Discussion** | Our analyses suggest children younger than 5 years with mild to moderate COVID-19 have high amounts of SARS-CoV-2 viral RNA in their nasopharynx compared with older children and adults. Our study is limited to detection of viral nucleic acid, rather than infectious virus, although SARS-CoV-2 pediatric studies reported a correlation between higher nucleic acid levels and the ability to culture infectious virus.<sup>5</sup> Thus, young children can potentially be important drivers of SARS-CoV-2 spread in the general population, as has been demonstrated with respiratory syncytial virus, where children with high viral loads are more likely to transmit.<sup>6</sup> Behavioral habits of young children and close quarters in school and day care settings raise concern for SARS-CoV-2 amplification in this population as public health restrictions are eased. In addition to public health implications, this population will be important for targeting immunization efforts as SARS-CoV-2 vaccines become available.

Taylor Heald-Sargent, MD, PhD  
William J. Muller, MD, PhD  
Xiaotian Zheng, MD, PhD  
Jason Rippe, JD  
Ami B. Patel, MD, MPH  
Larry K. Kocielek, MD, MSCI

**Author Affiliations:** Division of Infectious Diseases, Department of Pediatrics, Ann & Robert H. Lurie Children's Hospital, Chicago, Illinois (Heald-Sargent, Muller, Zheng, Rippe, Patel, Kocielek); Northwestern University Feinberg School of Medicine, Chicago, Illinois (Muller, Zheng, Patel, Kocielek).

**Accepted for Publication:** May 31, 2020.

**Corresponding Author:** Taylor Heald-Sargent, MD, PhD, Division of Infectious Diseases, Department of Pediatrics, Ann & Robert H. Lurie Children's Hospital, 225 E Chicago Ave, Box 20, Chicago, IL 60611 (thsargent@luriechildrens.org).

**Published Online:** July 30, 2020. doi:10.1001/jamapediatrics.2020.3651

**Author Contributions:** Drs Heald-Sargent and Kocielek had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

**Study concept and design:** Heald-Sargent, Muller, Kocielek.

**Acquisition, analysis, or interpretation of data:** All authors.

**Drafting of the manuscript:** Heald-Sargent, Muller, Kocielek.

**Critical revision of the manuscript for important intellectual content:** All authors.

**Statistical analysis:** Heald-Sargent, Kocielek.

**Administrative, technical, or material support:** All authors.

**Study supervision:** Muller, Patel, Kocielek.

**Conflict of Interest Disclosures:** Dr Muller has received grants from Ansun BioPharma, Astellas Pharma, AstraZeneca, Abbott Laboratories, Janssen Pharmaceuticals, Karius, Merck, Melinta Therapeutics, Roche, Tetrphase Pharmaceuticals, and Seqirus. Dr Patel has received grants from Aqua Pharmaceuticals. Dr Kocielek has received grants from Merck. No other disclosures were reported.

**Funding/Support:** Drs Heald-Sargent and Kocielek were supported by grants T32AI095207 and K23AI123525, respectively, from the National Institute of Allergy and Infectious Diseases.

**Role of the Funder/Sponsor:** The funder had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

- Lu X, Zhang L, Du H, et al; Chinese Pediatric Novel Coronavirus Study Team. SARS-CoV-2 infection in children. *N Engl J Med*. 2020;382(17):1663-1665. doi:10.1056/NEJMc2005073
- Paules C, Subbarao K. Influenza. *Lancet*. 2017;390(10095):697-708. doi:10.1016/S0140-6736(17)30129-0
- Cao Q, Chen YC, Chen CL, Chiu CH. SARS-CoV-2 infection in children: transmission dynamics and clinical characteristics. *J Formos Med Assoc*. 2020;119(3):670-673. doi:10.1016/j.jfma.2020.02.009
- Liu Y, Yan LM, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis*. 2020;20(6):656-657. doi:10.1016/S1473-3099(20)30232-2
- L'Huillier AG, Torriani G, Pigny F, Kaiser L, Eckerle I. Culture-competent SARS-CoV-2 in nasopharynx of symptomatic neonates, children, and adolescents. *Emerg Infect Dis*. 2020;26(10):26.
- Moreira LP, Watanabe ASA, Camargo CN, Melchior TB, Granato C, Bellei N. Respiratory syncytial virus evaluation among asymptomatic and symptomatic subjects in a university hospital in Sao Paulo, Brazil, in the period of 2009-2013. *Influenza Other Respir Viruses*. 2018;12(3):326-330. doi:10.1111/irv.12518

### Preventive Health Care Utilization Among Youths Who Have Run Away, Experienced Homelessness, or Been Stably Housed

Although it is well established that youths with unstable housing face increased health risks compared with peers with stable housing, there is considerable heterogeneity in youths' experiences of housing instability, with implications for health, clinical practice, and policy.<sup>1,2</sup> For instance, adolescents who are homeless with their families may have unique health needs compared with those who are unaccompanied,<sup>3</sup> and youths who run away may be at particularly high risk for poor health outcomes.<sup>4</sup> Given that annual preventive visits are recommended for all adolescents, they may represent an opportunity to explore housing status and address health needs of